



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 9/64, 15/11, 15/57	A2	(11) International Publication Number: WO 98/04686 (43) International Publication Date: 5 February 1998 (05.02.98)
(21) International Application Number: PCT/CA97/00535 (22) International Filing Date: 25 July 1997 (25.07.97) (30) Priority Data: 60/021,008 26 July 1996 (26.07.96) US 2,203,745 25 April 1997 (25.04.97) CA (71)(72) Applicants and Inventors: DAY, Robert [CA/CA]; 833 rue Pierre, Sainte-Dorothée, Québec H2W 3S6 (CA). SEIDAH, Nabil, G. [CA/CA]; Apartment 1412, 200 de Gaspé, Îles-des-Soeurs, Québec H3E 1S6 (CA). MARTEL, Rémi [CA/CA]; 4865 Lafontaine, Montréal, Québec H1V 1R7 (CA). CHRETIEN, Michel [CA/CA]; Apartment 1404, 1 Côte Sainte-Catherine, Montréal, Québec H2V 1Z8 (CA). REUDELHUBER, Tim [CA/CA]; 671 Warwick Drive, Baie d'Urfé, Québec H9X 2P4 (CA). LECLERC, Guy [CA/CA]; 327 Lorraine, Rosemère, Québec J7A 4K1 (CA). (74) Agents: DUBUC, Jean, H. et al.; Goudreau Gage Dubuc & Martineau Walker, 800 Place Victoria Tower, Suite 3400, P.O. Box 242, Montreal, Quebec H4Z 1E9 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PRO-PROTEIN CONVERTING ENZYME (57) Abstract <p>A cDNA clone encoding the human prohormone convertase PC5 was isolated from human adrenal gland messenger RNA. The deduced protein sequence would encode a 915 amino acid prepro PC5 which shares a very high degree of homology with the previously cloned rat and mouse homologues. PC5 mRNA is detected in multiple human tissues, including the brain, adrenal and thyroid glands, heart, placenta, lung and testes. PC5 mRNA was undetectable in the liver and is present at lower levels in skeletal muscle, kidney, pancreas, small intestine and stomach. Co-transfection of human PC5 and human prorenin expression vectors in cultured GH₄Cl cells leads to secretion of active renin. The activation of human prorenin by PC5 is dependent on a pair of basic amino acids at positions 42 and 43 of the prorenin prosegment and occurs only in cells containing dense core secretory granules. Human PC5 was co-localized with renin by immunohistochemistry in the zona glomerulosa of the adrenal gland suggesting that it could participate in the activation of a local renin-angiotensin system in the human adrenal cortex. PC5 is overexpressed in atherosclerotic coronary blood vessels. Silencing PC5 expression with a specific antisense oligonucleotide efficiently inhibited the proliferation of smooth muscle cells in culture. Furthermore, the antisense inhibited carotid stenosis in a carotid injury model. These results indicate that silencing PC5 applies to the prevention of restenosis. PCs could be targets of choice for treating any proliferative diseases involving their action on a given growth factor. Finally, the antisense oligonucleotide PC5 is to be used for silencing the activity of this enzyme towards HIV gp160, since both coexist in CD4⁺ T lymphocytes and the viral glycoprotein is cleavable by PC5.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE:

PRO-PROTEIN CONVERTING ENZYME

FIELD OF THE INVENTION:

This invention relates to protein processing enzymes or
5 pro-hormone convertases (PCs), specifically to PC5, more
specifically to the human PC5.

BACKGROUND OF THE INVENTION:

Pro-hormone convertases (PCs) belong to a family of
enzymes responsible for the maturation of proteic precursors
10 into active proteins or enzymes. Up to now, many human
enzymes of that family have been identified, namely furin,
PC1, PC2, PC4 and PC7. Each enzyme has a tissular
distribution which may be restricted (for example, PC4 is
restricted to male germ cells) or ubiquitous (furin is such
15 an example). Although all these enzymes share the properties
of cleaving precursor proteins at basic or dibasic residues,
they nevertheless have differing cleavage specificities. The
action of a specific pro-hormone convertase is therefore
governed by the cleavage sequence of a given protein
20 substrate, and/or by the location of that enzyme in a tissue
expressing or responding to a given proteic substrate growth
factor or hormone.

Renin is an aspartyl protease which makes an important
contribution to cardiovascular physiology and pathophysiology
25 through its key role in the synthesis of the vasoactive
octapeptide angiotensin II (AII). While the kidney is the
primary source of circulating active renin, several
additional tissues, including the pituitary and adrenal
glands, placenta, uterus, ovary, testes, heart, vasculature
30 and brain express the renin gene (reviewed in ¹⁻⁴). The
presence of additional components of the RAS (renin-
angiotensin system) in these tissues, including angiotensin
converting enzyme (ACE) and angiotensin II receptors, has led
to the proposal that certain tissues might contain a locally
35 active tissue renin-angiotensin system (tRAS) although the
actual function of the various tRAS is still largely a matter
of conjecture.

Renin is first synthesized as an enzymatically inactive precursor, prorenin, which is converted to active renin by the proteolytic removal of a 43 amino acid amino-terminal prosegment. The activity of the RAS within any given tissue would, therefore, be dependent on the existence of proteolytic enzymes capable of converting prorenin to active renin and on the expression of such prorenin processing enzymes (PPEs) in the same cells that express prorenin. The identity of the enzyme(s) responsible for the proteolytic activating human prorenin *in vivo* is still uncertain. Furthermore, it is possible that multiple PPEs exist in humans and these may differ among renin-producing tissues. Biochemical and microscopic studies of renin in the kidney suggest that candidate PPEs should be selective for cleavage of human prorenin at Lys⁴², Arg⁴³ of the prosegment⁵ and would be active in secretory granules of the juxtaglomerular (JG) cells.⁶ The lysosomal enzyme cathepsin B has been co-localized with human renin/prorenin in the secretory granules of JG cells and human pituitary lactotrophs^{7,8} and has been shown to cleave human prorenin *in vitro* with a high affinity and selectivity for the proper cleavage site.⁹ The prohormone convertase PC1 has also been shown to cleave human prorenin with the correct site-and organelle specificity in transfected cells¹⁰ and to co-localize with renin in the adrenal medulla and derived tumors¹¹, but not in JG cells.¹²

In an effort to identify novel PPEs, we recently determined the distribution of processing enzymes in an established renin-expressing tissue culture cell line derived from an oncogene-induced mouse tumor (As4.1 cells¹³). One such enzyme, the mouse prohormone convertase PC5, was found. Mouse PC5 is capable of partially cleaving human prorenin.

Miranda et al. (38) describe the cDNA and protein sequences for a human PC6 enzyme obtained by PCR from CD4⁺ T lymphocytes. PC5 and PC6 are different names given to what appears to be the same enzyme. However, the sequences of Miranda et al. comprise a plurality of substitutions when compared to the present PC5 sequences. Moreover, the size of messenger RNA encoding PC6 and PC5 are similar but not identical. Since the present PC5 sequences were obtained from

human adrenals, both enzymes may be isoforms, differentially expressed in tissues and they may have different activities.

Prorenin and HIV gp160 are most probably not the only proteic precursors to be recognized and cleaved by PC5. Many growth factors responsible for cell proliferation are cleaved by one or more PCs: they include platelet-derived-growth factors A and B (PDGF's), epidermal-growth-factor (EGF), insulin-like growth factors I and II (IGF's), transforming growth factors α and β (TGF's). Each of these named growth factors has the typical cleavage site motif K/R-(X)_n-R! (where n=0,2,4,6). Full biological potency is conferred to these growth factors only after cleavage at these sites, by one or more of the PC enzyme family. There is therefore a possibility that manipulating the expression of the PCs would affect cell proliferation via deficient growth factor activation.

Out of the >450,000 patients/year in the U.S. and Canada who undergo percutaneous transluminal coronary angioplasty (PTCA), 30-50% of them will restenose their coronaries within 3-6 months. This flare-up of endothelial and smooth-muscle cells proliferation is due to the activation of numerous regulatory growth factors. Therefore, knowing which enzyme(s) is (are) responsible for this activation, and manipulating the level of expression of this or theses enzyme(s) would be particularly useful to prevent restenosis.

STATEMENT OF THE INVENTION:

The present invention relates to the human PC5 (hPC5). We demonstrate that hPC5 isolated from human adrenals proteolytically activates human prorenin with the expected site- and organelle- specificity and that it is co-expressed with prorenin in the zona glomerulosa of the adrenal cortex. Therefore, PC5 is a prorenin-processing enzyme (PPE). Silencing the expression of PC5 would find a specific application in inhibiting the production of renin, and a method of inhibiting the production of renin is an object of the invention. Since the production of renin is one of targets of the RAS involved in hypertension. Furthermore, we demonstrate that hPC5 is overexpressed in atherosclerotic

coronary arteries. Antisense oligonucleotides have been designed, amongst which one has been shown to successfully silence the expression of hPC5 in smooth muscle cells in culture. This antisense inhibited carotid stenosis in a in vivo rabbit carotid injury model. These results indicate that a method of silencing the expression of PC5 would find a specific application in preventing restenosis.

PC5 is known to be expressed in CD4⁺ T cells, along with furin and PC7. The three enzymes are capable of converting HIV gp160 into its fusigenic form. Therefore, antisense constructs, particularly the oligonucleotide that successfully inhibited restenosis, will find a use in inhibiting expression of the activity of PC5 towards HIV gp160.

The complete amino acid and nucleotide sequence of hPC5 is described hereinbelow and are another object of this invention. Recombinant vectors and hosts comprising as a new insert, whole or part of hPC5, are also an object of the invention.

Oligopeptides derived from the proteic sequence of hPC5 are also an object of the invention.

Antibodies directed against the whole protein hPC5 or a part thereof are also an object of the invention.

Diagnostic methods and kits comprising oligonucleotides or antibodies binding PC5 nucleic acids or protein or peptides are also an object of the invention.

This invention will be described hereinbelow by way of specific embodiments, examples and figures which purpose is to illustrate the contemplated aspects of the invention, and not to limit the scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1: Schematic diagram of the isolated cDNAs encoding hPC5. Restriction enzyme sites used in sub-cloning are denoted. Solid lines represent clones isolated from a phage library. Hatched lines denote the portion of the cDNA isolated by RT-PCR of human adrenal mRNA. The double line represents the portion of the mouse PC5 cDNA (corresponding

to the amino terminus of the signal peptide) which was used to complete the cDNA for expression.

5 Figure 2: Nucleotide and derived protein sequence of hPC5
(SEQ ID NOS: 1 and 2, respectively). Proposed signal peptide
(solid arrow) and prosegment (open arrow) cleavage sites are
denoted based on data from mouse PC5.¹⁵ The underlined
sequence represents the portion of the signal peptide from
mouse PC5 which was used in the expression vector
construction.

10 Figure 3: Distribution on PC5 RNA in various human tissues.
Each lane contains 2 μ g poly-A RNA. Filters were hybridized
with a radiolabeled probe for hPC5 as described in Materials
and Methods. Shown at left is the migration of single strand
size standards in kilobases (Kb). Note that the absolute
15 signal cannot be compared between the two filters as they
were of different ages and hybridized at different times.

Figure 4: hPC5 cleaves human prorenin with site and cell
specificity. Panel A: GH₄C₁ cell were co-transfected with
expression vectors for the indicated proteins. Supernatants
20 were collected 30 hrs. after transfection and assayed for %
active renin [(active renin/total renin) X 100]. Bars
represent the mean \pm S.E.M of 9 independent transfections. *
= P < 0.0001 as compared to proren + pUC, as determined by the
Mann-Whitney non-parametric test. Panel B: Resulting
25 secretion of active renin after co-transfection of CHO cells
with an expression vector for prorenin and either a control
plasmid (pUC) or hPC5. Bars represent the mean \pm S.E.M of 3
independent transfections.

Figure 5: Active renin generation in secretory granules of
30 co-transfected GH₄C₁ cells. Parallel wells of GH₄C₁ cells co-
transfected with expression vectors for human prorenin and
human PC5 were incubated for 20 min. in medium containing
either 50mmol/L NaCl (control) or 50mmol/L KCl (a de-
polarizing agent which causes the acute release of secretory
35 granules). Percent active renin was calculated as described
in the legend to Figure 4A. Bars represent the mean \pm SEM of
3 independent transfections. * = P < 0.005 using Student's t-
test.

Figure 6: Immunodetection of hPC5 and renin/prorenin in renal cortex, human placental cotyledon and adrenal gland. Positively stained areas are denoted by solid arrows. Sections in adrenal cortex are separated by 5mM to show co-localization in the cells of the zona glomerulosa (g) and absence of staining in the capsule (c) and zona fasciculata (f). Original magnification 25X (kidney and placenta) and 80X (adrenal gland).

Figure 7 : In vivo hybridization analysis of PC5 mRNA in human coronary blood vessels in atherosclerosis. The lower panel show a vessel where a severe lesion was observed. PC5 mRNA was abundantly expressed in this vessel, in the smooth muscle cells in the neointimal formation (see arrows). In comparison, another vessel which is free of any lesion, did not express PC5 mRNA (shown in the upper panel).

Figure 8: Western blot analysis of PC5 protein in rabbit smooth muscle cells treated with either antisense, sense or mismatch PC5 oligonucleotides. The specific PC5 band is identified (see arrow) by comparison with proteins extracted from rat supraoptic nucleus (SON) and SK-N-MCIXC cells (human neuroepithelioma). In the sample extracted from antisense PC5 treatment, we observe a dramatic decrease in the level of PC5 signal (approximately 2-3 fold decrease) in comparison to the control sense or mismatch PC5 oligos. This indicates that the antisense treatment reduced significantly the protein levels of PC5 in rabbit smooth muscle cells.

Figure 9 : Rabbit in vivo test of the PC5 antisense ODN as compared to the control sense and random ODNs. It shows a decreased stenosis due to the presence of a PC5 antisense when compared to the sense and random controls.

Figure 10 : Proprotein convertase immunoreactivity in human atherectomy specimens. It shows the presence of the enzymes of the pro-hormone convertase family which are present in the specimens.

Figure 11: Illustrates differences between cDNA sequences of PC6 (Miranda et al. SEQ ID No. 4) and PC5 (present invention).

Figure 12: Illustrates differences between protein sequences of PC6 (Miranda et al. SEQ ID No. 5) and PC5 (present invention).

DESCRIPTION OF THE INVENTION

5 Materials and Methods

cDNA library construction and screening: A cDNA library derived from total human adrenal RNA was constructed by Stratagene (La Jolla, CA) in the phage vector Uni-Zap XR. Six hundred thousand phage plaques were screened initially using
10 radioactive probes and standard methodologies.¹⁴ The initial hybridization probe was a 320 base pair DNA fragment derived from reverse-transcriptase PCR of human brain RNA using information derived from an unidentified human cDNA sequence tag in Genbank (Accession # M85522) with a high degree of
15 similarity to the previously cloned mouse PC5.¹⁵ Fragment labeling was carried out using ³²P dCTP and a random primer labeling kit (Boehringer-Mannheim Canada, Laval, Quebec, Canada) according to manufacturer's instructions. One positive hybridizing phage (hPC5A) was identified. Its insert
20 was sequenced in its entirety using the dideoxy-chain termination method and found to code for an 1150 base pair cDNA with a high degree of sequence similarity to mouse PC5 (data not shown). A 1070 base pair fragment (excluding the poly A tail) was excised from hPC5A, labeled and used to re-
25 screen an additional 600,000 phage from the cDNA library. A second phage clone (hPC5B) was isolated and found to contain an 1807 base pair cDNA insert overlapping hPC5A and extending toward the 5' end of the cDNA (Figure 1).

Reverse-transcriptase PCR: One microgram of poly A+ RNA
30 from total human adrenal (Clontech Laboratories, Palo Alto, CA) was subjected to reverse transcriptase polymerase chain reaction (RT-PCR) using a published procedure¹⁶ and the following oligonucleotides:

Forward oligonucleotide; derived from a region corresponding
35 to the signal peptide of mouse PC5.¹⁵ An artificial HindIII restriction enzyme cleavage site added to the 5'-end of the amplified

fragment for the purpose of cloning is underlined:

5'-CCAAGCTTGGCTGCTGTGCGTGCTGGC-3'

Reverse oligonucleotide; derived from the 5'-end of the phage hPC5B. An internal BglIII restriction enzyme site is underlined: 5'-CTGCCTCAGATCTGTAGTG-3'

5 The entire RT-PCR reaction was repeated 4 times and 4 independently derived clones of the amplified fragment were sequenced and the sequences were compared. The sequence submitted to Genbank (Accession #U49114) represents the consensus sequence, defined as any nucleotide appearing in
10 3/4 clones.

Northern blot analysis: Tissue distribution of PC5 mRNA was determined by hybridizing commercially purchased nitrocellulose filters containing aliquots (2µg) of Poly-A RNA from various human tissues (Clontech Laboratories, Palo
15 Alto, CA). The probe used was a complementary RNA derived from the full length hPC5 cDNA. Probe labeling and hybridization were carried out as previously described.¹⁷

Expression vector construction: A cDNA fragment from the KpnI site (Figure 1) to just past the stop codon was excised
20 from the phage hPC5B and combined with a KpnI to HindIII (see above) fragment derived from portions of two independent RT-PCR clones (so as to eliminate errors arising from the Taq polymerase). A region corresponding to the first 16 amino acids of the signal peptide derived from mPC5 was attached to
25 the 5'-end by overlap-extension PCR.¹⁸ Thus, the entire cDNA, encoding amino acids 1-16 derived from the mPC5 signal peptide and the remainder from hPC5, was subcloned into the expression vector RSV-globin¹⁹ which places the cDNA under the control of the RSV promoter and provides a 3' intron and
30 polyadenylation signal from the rabbit beta globin gene. The entire subcloned fragment was subsequently verified by DNA sequencing.

Cell culture and transfection: GH₄C1 cells were plated in 6-well culture dishes at a density of 5X10⁵ cells per
35 well. Twenty four hours later, medium was changed and the cells were transfected by the DEAE-dextran method using a commercial kit (CellPfect Transfection kit, Pharmacia Biotech, Baie D'urfe, Quebec, Canada) according to manufacturer's instructions. Each well received 0.18 µg of

either the hPC5 expression vector or a neutral plasmid vector (pUC18) and 0.18 μ g of an expression vector for human prorenin (pRHR1100) or its equivalents in which amino acids 42 or 43 of the prorenin prosegment were mutated to alanine (K/A -2 and R/A-1, respectively²⁰). Supernatants were collected 30 hrs. after transfection and assayed for prorenin and renin content as previously described.²⁰

To verify that conversion of the prorenin occurred in the secretory granules, GH₄Cl transfected with the human prorenin and hPC5 expression vectors were stimulated to release secretory granules by depolarization using a previously published technique.²¹ Forty hrs. after co-transfection, the culture medium in parallel wells of transfected cells was replaced with pre-warmed medium supplemented to a final concentration of 50 mmol/L with either NaCl (control) or KCl (secretagogue). The media were collected after 20 min. and assayed for renin/prorenin. A potassium-dependent increase in the percent active renin contained in cell supernatants was taken as an indication of active renin release from the secretory granules of the transfected cells. Results shown in Figure 5 represent the mean of three independent transfection experiments.

Immunolocalization of hPC5 in human tissues: Human tissue was obtained post-mortem (kidney and adrenal gland) or post-partum (placental cotyledon), fixed in Bouin's solution and embedded in paraffin. For immunolocalization, 5 μ m sections were mounted on gelatin-coated slides, deparaffinized and incubated with a 1:50 dilution of a polyclonal rabbit antiserum raised against a peptide corresponding to the N-terminal 16 amino acids of rat PC5 (PC5.MAP antibody) or a 1:200 dilution of a polyclonal rabbit antiserum against recombinant human prorenin. For kidney and placental specimens, immune complexes were revealed by incubation with protein A-colloidal gold (15 nm particles) synthesized from tetra-chloroauric acid (BDH) according to the method of Ghitescu and Bendayan.²² Gold particles were enhanced for viewing in the light microscope by incubation with silver (IntenseTM Silver Enhancement Kit, Amersham Life Science, Oakville, Ontario, Canada) and sections were

counter-stained with hematoxylin and methyl green. Immune complexes on human adrenal sections were detected with a 1:200 dilution of biotin-labeled donkey anti-rabbit IgG and a 1:300 dilution of streptavidin-horseradish peroxidase complex (Amersham Life Science, Oakville, Ontario, Canada) and were incubated with diaminobenzidine and hydrogen peroxide (Sigma Chemicals, St. Louis, MO) as chromogen. All positive staining patterns were subsequently verified for specificity by omission of the first antibody.

10 Results

The primary sequence of human PC5 is shown in Figure 2. We were unable to clone the extreme 5'-end of the cDNA either by the RACE protocol¹⁶ or by using oligonucleotides based on the published sequence of mouse PC5^{15,23}, possibly due to a high G/C content of the cDNA in this region. However, based on the published cDNA sequences for rat and mouse PC5¹⁵, we are confident that we have isolated all but the 5'-most portion of the cDNA corresponding to the first 12 amino acids of the signal peptide. By comparison with the published sequence of mouse PC5, we predict that the cDNA isolated would code for a preproPC5 of 915 amino acids, including a signal peptide and a prosegment of 32 and 84 amino acids, respectively. The deduced sequence of hPC5 is 88% identical to the previously published mouse PC5 cDNA and 96% identical to the mouse PC5 protein.

Northern analysis of poly A RNA from a variety of human tissues reveals a major band of approximately 6.6 Kb and a minor band at approximately 3.8 Kb (Figure 3). PC5 RNA is detected in the brain, heart, placenta, lung, thyroid gland and testes and at lower levels in the skeletal muscle, kidney and pancreas, small intestine and stomach. In the adrenal gland, PC5 is particularly enriched in the cortex (Figure 3).

Because PC5 RNA appears to be expressed in a number of tissues previously reported to contain active renin, we have tested the ability of hPC5 to cleave human prorenin in a cell co-transfection assay (Figure 4A). As has been previously reported¹⁰, when cultured rat sommatotrophic GH₄C1 cells are co-transfected with an expression vector encoding human

prorenin and a neutral plasmid vector, only unprocessed prorenin is secreted into the culture supernatant. In contrast, if the human prorenin expression vector is co-transfected with an expression vector encoding human PC5, a portion of the expressed prorenin is secreted as active renin. Co-expression of human PC5 with prorenin mutated at either of the basic residues forming the native cleavage site (Lysine 42 or Arginine 43) prevents activation. These results suggest that human PC5 activates human prorenin by proteolytic cleavage at the site previously reported for activation of renin in humans.⁵ While human PC5 cleaves human prorenin in GH₄C1 cells, there is no apparent increase in active renin secretion when co-transfections are carried out in Chinese Hamster Ovary (CHO) cells (Figure 4B). One obvious difference in the CHO cell line as compared to GH₄C1 cells is their lack of secretory granules, suggesting that either human PC5 or human prorenin or both require the secretory granule environment for this proteolytic step. This conclusion is supported by the acute increase in active renin detected in the supernatants of co-transfected GH₄C1 cells treated for 20 min. with potassium chloride (Figure 5), a depolarizing agent which causes the release of secretory granules.²¹

Using a polyclonal antibody raised against a peptide derived from mouse PC5, we have studied the distribution of human PC5 in several human tissues (Figure 6). To date, we have been unable to detect staining for PC5 in the human kidney, although our sections stain positively for renin. In the placental cotyledon, PC5 is located in the syncytiotrophoblast layer of the chorionic villi while anti-renin antibody stains primarily the chorionic mesoderm. In the adrenal gland, the antibodies against both renin and PC5 show a preferential staining of zona glomerulosa cells in the adrenal cortex (g) with very little staining of the capsule (c) and zona fasciculata (f). No staining was evident with omission of the first antibody (data not shown). Thus, our immunohistochemical studies would suggest that, of the three tissues studied, it is likely that prorenin and PC5 are only

clearly co-localized in the zona glomerulosa of the human adrenal cortex.

Discussion

In the present study, we describe the cloning and
5 expression of the human prohormone convertase PC5 and its
activity as a human PPE. Co-transfection assays in cultured
cells demonstrates that hPC5 activates human prorenin with
the expected site-specificity and that this cleavage most
likely takes place in dense core secretory granules. In
10 addition, immunohistochemistry of human tissues shows co-
localization of hPC5 with renin in the zona glomerulosa of
the adrenal cortex.

Several lines of evidence suggest that the human adrenal
gland contains a physiologically important local RAS: First,
15 RNA encoding angiotensinogen and renin have been detected in
preparations from the human adrenal zona glomerulosa,
fasciculata and medulla^{24,25}, confirming that both renin and
its substrate are synthesized within the human adrenal gland.
Second, ACE inhibition or blockade of angiotensin receptors
20 inhibits aldosterone release from human adrenal tissue
explants²⁶, suggesting that the local RAS plays an active role
in the regulation of aldosterone secretion from the adrenal
gland. Third, tissue explants of human adrenal cortex and
aldosterone-secreting adenomas secrete small quantities of
25 active renin^{24,26,27}, suggesting that the adrenal cortex
expresses a PPE capable of activating human prorenin. Our
current results suggest that PC5 could be the PPE responsible
for activation of renin in the human adrenal cortex as both
renin and hPC5 are immuno-detectable in the zona glomerulosa.
30 Additional circumstantial evidence supports this conclusion:
First, centrifugal fractionation of adrenal cortical cells
reveals that renin is contained in the "granular" fraction,
of intermediate density between vesicles and lysosomes.²⁸ As
our current study suggests that PC5 only cleaves human
35 prorenin in cells containing secretory granules, renin would
be in the appropriate intracellular compartment to be
activated by PC5 in the adrenal cortex. Second, rats
transgenic for mouse Ren-2 renin [TGR(mRen-2)27] display

fulminant hypertension²⁹ which correlates best with the expression of the mouse prorenin in the adrenal gland.³⁰⁻³² As previous studies have demonstrated that PC5 is capable of activating mouse Ren-2 prorenin, but not rat prorenin (²³and data not shown) it is possible that the TGR(mRen-2)²⁷ transgenic rat is a model for activation of a tissue RAS by the fortuitous juxtaposition of prorenin with an appropriate PPE in the adrenal cortex. These results also raise the possibility that the tissue-distribution of PPEs and their apparent selectivity in activating prorenin from different species could lead to differing functions of the tissue RAS between rodents and humans.

The principal source of circulating active renin in humans is the JG cells of the kidney. Although low levels of hPC5 RNA can be detected by Northern blot analysis in a sample of total kidney poly-A RNA (Fig. 3), we were unable to localize PC5 immunostaining in kidney sections (Fig. 6) raising the possibility that PC5 is expressed at low levels in diffuse cell types in the kidney. Thus, while these results do not formally rule out PC5 as a PPE in the kidney, our inability to detect it in JG cells makes it unlikely that it plays a major role in the production of renal renin. In contrast, relatively abundant amounts of PC5 mRNA and protein were detected in the placenta although evidence suggests that placental cells in culture³³ and *in vivo*³⁴ only secrete prorenin. However, immunostaining revealed that the cells producing PC5 and prorenin in the human placenta are distinct. It is also unlikely that PC5 would activate prorenin once the two proteins are secreted due to the apparent requirement of a granular environment for the cleavage of prorenin by hPC5 in transfected cells. Thus, in contrast to the case in the adrenal gland, it is unlikely that PC5 expressed in the human placenta would activate placental prorenin.

In the mouse, two forms of hPC5 have been predicted based on cloned cDNAs; the first would be analogous to the hPC5 cDNA described in this study and to that cloned from rat tissues^{15,23} while the second, called PC6B, is extended at its 3'-end due to a differential RNA splicing event.³⁵ Although

the hPC5 cDNA we have cloned is only roughly 3 Kb in length, the major RNA band seen in human tissues is of approximately 6.6 Kb. The identity of the longer band hybridizing to the hPC5 probe is currently unknown. It should be noted that
5 neither of the cDNA clones isolated from a screening of 1.2 million phage from the adrenal library was extended at its 3'-end (Figure 1), although the probes used in their isolation cover the region of homology with the mouse PC6B variant.³⁵ In mouse tissues, expression of the PC6B variant is
10 restricted to few tissues³⁵ while the abundance of the 6.6Kb detected with the hPC5 probe is directly proportional to the abundance of the 3.8Kb band. Hybridization of RNA blots from rodent tissues using a PC5 probe also reveals RNA bands of 3.8, 6.5 and 7.5 Kb^{15,35} and use of a PC5-specific probe
15 reveals a band at 6.5 Kb. Thus, it is possible that additional PC5 RNA species exist in mammals that are extended at their 5'-ends. Alternatively, human tissues may be particularly enriched in a homologue to PC6B which was not picked up in our screenings. Recent data suggest that the
20 alternate C-terminal tail present on PC6B may serve to retain the enzyme in the Golgi network, while the "short" form of mouse PC5 is targeted to dense core secretory granules (N.G. Seidah, unpublished). These data and the results of our co-transfection assays (Fig. 4) would suggest that the "short"
25 form of hPC5 described here is the form which would be active in renin processing in secretory granules.

The PC5 enzymes isolated from humans and mice show a remarkably high degree of conservation at the nucleotide and protein sequence levels. This degree of similarity is higher
30 than that seen for the other mammalian PC enzymes which seem to diverge in the C-terminal half of the enzyme.^{36,37} This high degree of sequence conservation may reflect an essential function of PC5 (and the C-terminus of PC5) in mammals.

PC5 is linked to smooth muscle proliferation

35 To investigate which PC could be a potential target of smooth muscle cell proliferation, we tested if any of the PCs were affected in the process of restenosis, wherein such proliferation is observed. Changes in PC levels in the

process of restenosis is a distinct possibility since in previous studies using animal models or cell lines, we have shown that PC levels can be regulated or even be induced. We thus obtained human restenosed coronary tissues from patients. These tissues were screened for each of the PC mRNAs using *in vivo* hybridization histochemistry in order to obtain information within an anatomical context. Coronaries with partial or total occlusions demonstrated dramatically increased PC5 mRNA levels within smooth muscle tissues, whereas coronary tissue without occlusions were PC5 negative. These results indicate that PC5 is either strongly up-regulated or induced in the human coronary arteries during the active process of stenosis (fig. 7). To our knowledge this is the first indication that a specific PC is directly linked to smooth muscle proliferation.

These results suggested that if PC5 enzymatic activity could somehow be inhibited or the upregulation of PC5 mRNA could be prevented, this may attenuate or stop the process of restenosis. This could occur through the inhibition of the processing function of this enzyme on the numerous growth factors that are involved in the formation of the coronary lesion. If these growth factors are not processed they will remain biologically inactive. Our approach was to test the effectiveness of PC5 antisense inhibition on smooth muscle proliferation *in vitro*.

A specific antisense oligonucleotide (ODN) was shown to drastically inhibit smooth muscle proliferation using an *in vitro* model of rabbit smooth muscle in culture. Incubating rabbit smooth muscle cells with a PC5 antisense 17-mer oligonucleotide shown in Table 1 caused a dose-dependent inhibition smooth muscle proliferation with a maximal inhibitory effect of 81.6% + 1.6% at 10 mM (mean of three experiments done in quadruplicates). This inhibitory effect is highly significant ($P=0.0001$) as compared to controls which included either a sense or a mismatched oligonucleotide used at the same concentration (see Table 1). In addition we found that the expression of PC5 is decreased in the affected cells (fig. 8). When compared to other targets, such as *c-myc*, this approach was much more effective in inhibiting

smooth muscle proliferation, as the best effects of antisense *c-myc* resulted in $71.7 \pm 3.5\%$ (means of three experiment done in quadruplicate) inhibition (mean of three experiment done in quadruplicates). These results are indicative of an *in vivo* effect since silencing PC5 would impede muscle cell proliferation and restenosis.

Cholesterol conjugation of oligonucleotides

Phosphorothioate antisense ODNs were synthesized on a DNA/RNA synthesizer following standard procedure (Applied Biosystems). Conjugation of oligomers with cholesterol was achieved with 3'-cholesterol-VN CPG (Clontech), a virtual nucleotide (VN) glass reagent that introduces a cholesterol label to the 3' terminus of an oligonucleotide via solid-phase synthesis. When ODN synthesis was completed, oligomers were removed from the column with 30% NH_4OH (1 hour at room temperature), and then deprotected for 8 hours at 60°C . Oligos were purified and detritylated with oligonucleotide purification cartridges (Applied Biosystems), and then lyophilized with a centrifugal evaporator (Savant SpeedVac).

In vivo arterial ODN transfection

New Zealand rabbits male or female (2 Kg) were intramuscularly sedated with xylazine (2 mg/Kg) and anesthetized with ketamine (100 mg/Kg) prior to surgical exposure of left carotid artery. Segments of 10 mm of carotids were transiently isolated by temporary ligatures and rinsed with 0.9% sodium chloride via a cannula until there was no more visible evidence of blood components. Carotid arteries were transfected with $80 \mu\text{mol/L}$ of antisense ODNs in a 1 cm portion either alone or conjugated to cholesterol for a period of 30 minutes. The volume infused was $100 \mu\text{l}$, and no visible loss of volume was noted throughout the incubation period. Following transfection, the treated segments were rinsed with 0.9% sodium chloride ($3 \times 100 \mu\text{l}$) and upon cannula removal, the arteriotomy site was repaired with microsutures, restoring normal blood flow and the neck wound closed. All experimental protocols in this project were

approved by the Institutional Committee for Animal Protection of the Louis-Charles Simard Research Center.

Neointimal hyperplasia inhibition

A total of 36 New Zealand white rabbit carotid arteries were injured with a 2.5 mm balloon catheter serially inflated for 1 minute to 4, 6, 8 and 10 atm with gentle traction, allowing 45 seconds between inflations. Two weeks later, a second injury was imposed at the same arterial site which was then transfected in a 1 cm portion with 80 $\mu\text{mol/L}$ (100 μl of volume injected) of therapeutic molecules or with 100 μL of NaCl 0.9% as control. Intimal/medial areas were evaluated by computer analysis on histological sections derived from transfected arteries two weeks following the second injury and transfection procedure.

The addition of a PC5 antisense 17-mer ODN shown in Table 1 at the time of a second carotid injury with a balloon catheter decreased carotid stenosis, measured as area ratio intima/media, by 40% (Area ratio intima/media sense ODN minus area ratio antisense ODN divided by area ratio sense ODN; see figure 9). This inhibitory effect is highly significant ($P = 0.0118$ and $P = 0.0078$) as compared to the sense and random controls, respectively. These results are the basis of a method of preventing stenosis comprising administering an effective stenosis inhibitory dose of a PC5 antisense to a subject in need for such a treatment. Other antisense oligonucleotides may be added to optimize this method of preventing restenosis, such as those silencing the expression of other convertases, namely PC2, which are also observed in atherectomy specimens (see Figure 10).

The development of drugs based on the inhibition or the inactivation of the convertases is of great interest because the drugs can easily be delivered directly at the affected site during the intervention by the cardiologist.

In addition we claim that this therapeutic approach, based on the inhibition of cell growth by antisense against one of the convertases will be applicable to all

proliferative diseases involving maturation of a given proteic precursor into an active protein.

TABLE 1

The sequences of the oligonucleotides used are:

5	Antisense	GCAACTTGCCAGAGCAT	SEQ ID NO: 3
	Sense	ATGCTCTGGCAAGTTGC	
	Random	AATCCGTGAGACCAGTC.	

PC5 is involved in the cleavage of HIV gp160 into gp120 and gp41.

10 As mentioned above, PC5, PC7 and furin are known to be present in CD4⁺ T lymphocytes. All three enzymes cleave HIV gp160 to gp120 and gp41 as well as a synthetic peptide covering the junction wherein cleavage occurs in gp160. Since the 17-mer antisense ODN defined in Seq ID. No. 3
15 successfully silenced the expression of PC5 and prevented restenosis, the same oligonucleotide as well as any other oligonucleotide or construct having an equivalent silencing function will find use in inhibiting the action of PC5 on HIV gp160 in CD4⁺ T lymphocytes. To optimize the inhibition of
20 conversion of gp160 into its fusiogenic form, a cocktail comprising antisense molecules to PC5, PC7 and furin is contemplated as part of the present invention. Appropriate vehicles such as liposomes may be used to deliver these antisense molecules to the target tissues.

25 This invention has been described hereinbelow. It may be apparent to the skilled reader that modifications can be made thereto without departing from the above teachings. These modifications are considered as part of the scope of the present invention, as defined in the appended claims.

References

1. Frohlich ED, Iwata T, Sasaki O. Clinical and physiologic significance of local tissue renin-angiotensin systems. *Am J Med.* 1990;87 (suppl 6B):6B-19S-6B-23S
- 5 2. Johnston CI, Burrell LM, Perich R, Jandeleit K, Jackson B. The tissue renin-angiotensin system and its functional role. *Clin Exper Pharmacol Physiol.* 1992;19:1-5
3. Gardner DG, Gertz BJ, Deschepper CF, Kim DY. Gene for the rat atrial natriuretic peptide is regulated by glucocorticoids *in vitro.* *J Clin Invest.* 1988;82:1275-1281
- 10 4. Lee MA, Bohm M, Paul M, Ganten D. Tissue renin-angiotensin systems. Their role in cardiovascular disease. *Circulation.* 1993;87:IV7-113
5. Do YS, Shinagawa T, Tam H, Inagami T, Hsueh WA. Characterization of pure human renal renin. *J Biol Chem.* 1987;262:1037-1043
- 15 6. Taugner R, Kim SJ, Murakami K, Waldherr R. The fate of prorenin during granulopoiesis in epithelioid cells. Immunocytochemical experiments with antisera against renin and different portions of the renin prosegment. *Histochemistry.* 1987;86:249-253
- 20 7. Taugner R, Hackenthal E. On the character of the secretory granules in juxtaglomerular epithelioid cells. *Int Rev Cytol.* 1988;110:93-131
- 25 8. Saint-Andre JP, Rohmer V, Pinet F, Rousselet MC, Bigorgne JC, Corvol P. Renin and cathepsin B in human pituitary lactotroph cells. An ultrastructural study. *Histochemistry.* 1989;91:291-297
9. Wang PH, Do YS, Macaulay L, Shinagawa T, Anderson PW, Baxter JD, Hsueh WA. Identification of renal cathepsin B as a human prorenin-processing enzyme. *J Biol Chem.* 1991;266:12633-12638
- 30 10. Benjannet S, Reudelhuber T, Mercure C, Rondeau N, Chretien M, Seidah NG. Proprotein conversion is determined by a multiplicity of factors including convertase processing, substrate specificity, and intracellular environment. Cell type-specific processing of human prorenin by the convertase PC1. *J Biol Chem.* 1992;267:11417-11423
- 35

11. Konoshita T, Gasc JM, Villard E, Takeda R, Seidah NG, Corvol P, Pinet F. Expression of PC2 and PC1/PC3 in human pheochromocytomas. *Mol Cell Endocrinol*. 1994;99:307-314
12. Reudelhuber TL, Ramla D, Chiu L, Mercure C, Seidah NG.
5 Proteolytic processing of human prorenin in renal and non-renal tissues. *Kidney Int*. 1994;46:1522-1524
13. Sigmund CD, Okuyama K, Ingelfinger J, Jones CA, Mullins JJ, Kane C, Kim U, Wu CZ, Kenny L, Rustum Y, Dzau VJ, Gross KW. Isolation and characterization of renin-expressing cell
10 lines from transgenic mice containing a renin-promoter viral oncogene fusion construct. *J Biol Chem*. 1990;265:19916-19922
14. Sambrook J, Fritsch EF, Maniatis T. *Molecular cloning: A laboratory manual*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press, 1989;
- 15 15. Lusson J, Vieau D, Hamelin J, Day R, Chretien M, Seidah NG. cDNA structure of the mouse and rat subtilisin/kexin-like PC5: a candidate proprotein convertase expressed in endocrine and nonendocrine cells. *Proc Natl Acad Sci USA*. 1993;90:6691-6695
- 20 16. *Current Protocols in Molecular Biology*. New York: John Wiley and Sons, Inc. 1996;
17. Day R, Schafer MK, Watson SJ, Chretien M, Seidah NG. Distribution and regulation of the prohormone convertases PC1 and PC2 in the rat pituitary. *Molec Endocrinol*. 1992;6:485-
25 497
18. Horton RM, Hunt HD, Ho SN, Pullen JK, Pease LR. Engineering hybrid genes without the use of restriction enzymes: gene splicing by overlap extension. *Gene*. 1989;77:61-68
- 30 19. Chu WN, Baxter JD, Reudelhuber TL. A targeting sequence for dense core secretory granules resides in the protein moiety of human prorenin. *Molec Endocrinol*. 1990;4:1905-1913
20. Chu WN, Mercure C, Baxter JD, Reudelhuber TL. Molecular determinants of human prorenin processing. *Hypertension*.
35 1992;20:782-787
21. Scammell JG, Burrage TG, Dannies PS. Hormonal induction of secretory granules in a pituitary tumor cell line. *Endocrinology*. 1986;119:1543-1548

22. Ghitescu L, Bendayan M. Immunolabeling efficiency of protein A-gold complexes. *J Histochem Cytochem*. 1990;38:1523-1530
23. Nakagawa T, Hosaka M, Torii S, Watanabe T, Murakami K, Nakayama K. Identification and functional expression of a new member of the mammalian Kex2-like processing endoprotease family: Its striking structural similarity to PACE-4. *J Biochem*. 1993;113:132-135
24. Racz K, Pinet F, Gasc JM, Guyene TT, Corvol P. Coexpression of renin, angiotensinogen, and their messenger ribonucleic acids in adrenal tissues. *J Clin Endocrinol Metab*. 1992;75:730-737
25. Paul M, Wagner J, Dzau VJ. Gene expression of the renin-angiotensin system in human tissues. Quantitative analysis by the polymerase chain reaction. *J Clin Invest*. 1993;91:2058-2064
26. Fallo F, Pistorello M, Pedini F, D'Agostino D, Mantero F, Boscaro M. in vitro evidence for local generation of renin and angiotensin II/III immunoreactivity by the human adrenal gland. *Acta Endocrinologica*. 1991;125:319-330
27. Sarzani R, Fallo F, Dessi-Fulgheri P, Pistorello M, Lanari A, Paci VM, Mantero F, Rappelli A. Local renin-angiotensin system in human adrenals and aldosteronomas. *Hypertension*. 1992;19:702-707
28. Mizuno K, Hoffman LH, McKenzie JC, Inagami T. Presence of renin secretory granules in rat adrenal gland and stimulation of renin secretion by angiotensin II but not by adrenocorticotropin. *J Clin Invest*. 1988;82:1007-1016
29. Mullins JJ, Peters J, Ganten D. Fulminant hypertension in transgenic rats harbouring the mouse Ren-2 gene. *Nature*. 1990;344:541-544
30. Yamaguchi T, Tokita Y, Franco-Saenz R, Mulrow PJ, Peters J, Ganten D. Zonal distribution and regulation of adrenal renin in a transgenic model of hypertension in the rat. *Endocrinology*. 1992;131:1955-1962
31. Rocco S, Rebuffat P, Cimolato M, Opocher G, Peters J, Mazzocchi G, Ganten D, Mantero F, Nussdorfer GG. Zona glomerulosa of the adrenal gland in a transgenic strain of

- rat: a morphologic and functional study. *Cell & Tiss Res.* 1994;278:21-28
32. Tokita Y, Franco-Saenz R, Mulrow PJ, Ganten D. Effects of nephrectomy and adrenalectomy on the renin-angiotensin system of transgenic rats TGR(mRen2)27. *Endocrinology.* 5 1994;134:253-257
33. Duncan KG, Haidar MA, Baxter JD, Reudelhuber TL. Regulation of human renin expression in chorion cell primary cultures. *Proc Natl Acad Sci USA.* 1990;87:7588-7592
- 10 34. Lenz T, James GD, Laragh JH, Sealey JE. Prorenin secretion from human placenta perfused in vitro. *Am J Physiol.* 1991;260:E876-E882
35. Nakagawa T, Murakami K, Nakayama K. Identification of an isoform with an extremely large Cys-rich region of PC6, a 15 Kex2-like processing endoprotease. *FEBS Lett.* 1993;327:165-171
36. Seidah NG, Hamelin J, Gaspar AM, Day R, Chretien M. The cDNA sequence of the human pro-hormone and pro-protein convertase PC1. *DNA & Cell Biology.* 1992;11:283-289
- 20 37. Seidah NG, Day R, Marcinkiewicz M, Chretien M. Mammalian paired basic amino acid convertases of prohormones and proproteins. *Ann NY Acad Sci.* 1993;680:135-146
38. Miranda L, Wolf J, Pichuantes S, Duke R, Franzusoff A. Isolation of the human PC6 gene encoding the putative host 25 proteas for HIV-1 gp160 processing in CD4⁺ T lymphocytes. *Proc. Natl. Acad. Sci. USA* 1996;93:7695-7700.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A purified human pro-protein converting enzyme named PC5.
- 5 2. A purified human pro-protein converting enzyme named PC5 isolated from adrenals.
3. A pro-protein converting enzyme as defined in claim 2 which has the amino acid sequence of SEQ ID NO: 2.
4. An isolated nucleic acid encoding the pro-protein
10 converting enzyme defined in claim 1, 2 or 3.
5. An isolated nucleic acid as defined in claim 4 which has the nucleic acid sequence of SEQ ID NO: 1.
6. An oligonucleotide having at least 15 nucleotides which is a part of the nucleic defined in claim 4 or 5 or its
15 complementary sequence.
7. An oligonucleotide which has the nucleic acid sequence defined in SEQ ID NO: 3.
8. A method of converting a proteic precursor cleavable by PC5 into a mature protein, which comprises the
20 steps of contacting said protein precursor with the pro-protein converting enzyme of claim 1 2 or 3 in conditions supporting the activity of said enzyme and recovering said protein.
9. A method of inhibiting muscle cell proliferation
25 which comprises the steps of contacting muscle cells with the oligonucleotide of claim 7 or any equivalent silencing antisense molecule.
10. A method as defined in claim 9, which results in the prevention of restenosis.
- 30 11. A method of silencing the expression of PC5 in a cell, which comprises the step of contacting said cell with the complementary sequence of the nucleic acid defined in claim 4 or 5, or a part thereof having at least 15 nucleotides.
- 35 12. A method as defined in claim 8, wherein said precursor is HIV gp160.
13. A method as defined in claim 11, wherein said cell is CD4⁺ T lymphocyte.

- 24 -

14. A method as defined in claim 13, wherein said cell is HIV infected T lymphocyte.


15. A method as defined in claim 14, which results in inhibition of cleavage of HIV gp160 by PC5.

5 16. A method as defined in any one of claims 11, and 13 to 15, wherein said part of said complementary sequence comprises the oligonucleotide defined in SEQ ID NO: 3.

10 17. A method as defined in claim 15, wherein said part of said complementary sequence comprises the oligonucleotide defined in SEQ ID NO: 3.

18. A method as defined in claim 17, further comprising antisense silencing molecules to PC7 and furin.

1/26

FIG - 1

ATGGACTGGGAAACCGCTGCAGCCGCGCGGACGGCGGGATCTGCTGTGCTGCTGCGGGGCTGCTCGCGGCTGTGCGGACGGCGCTACACCAACAC 120
M D W D W G N R C S R P G R R D L L C V L A L L G G C L L P V C R T R V Y T N H

TGGCAGTCAAAATGCGCGGGCTCCGGAGGCCAACCGTATCGCCAGCAAGTACGGATTTCATCAACATAGGACAGATAGGGCCCTGAAGGACTACTACCACTTCTACCATAGCAGG 240
W A V K I A G G F P E A N R I A S K Y G F I N I G Q I G A L K D Y Y H F Y H S R

ACGATTAAAGGTCAATTCTCGAGCAGAGGGACCCACAGTTTCAATTTCAATGGAACCAAGGTGGATCCAAACAGCAAGTGTGTAATAAAGCGGACAAAGAGGGATTATGACTTC 360
T I K R S V I S S R G T H S F I S M E P K V E W I Q Q V V K K R T K R D Y D F

AGTGTGCCAGTCTACCTATTTCAATGATCCCAAGTGGCCAGCATGTGATATGCACTGCAGTGACAATACACATCCCTGCCAGTCTGACATGAATATCGAAGGAGCCTGGAAGAGA 480
S R A Q S T Y F N D P K W P S M W Y M H C S D N T H P C Q S D M N I E G A W K R

GGTACACGGGAAGAACATTGTGTCACTATCTGTGATGCGGAATTGAGAGAACCCATCCAGATCTGATGCAAAACTACCATGCTCTGGCAAGTTGCGACGTGAATGGGAATGACTTG 600
G Y T G K N I V V T I L D D G I E R T H P D L M Q N Y D A L A S C D V N G N D L

GACCAATGCCTGTTATGATGCAAGCAAGAGCAAGCATGGGACTCGCTGTCTGGAGAAGTGGCAGCGCTGCAACAAATTCGCACTGCACAGTGGAAATTCCTTCAACGCCCAAG 720
D P H P R Y D A S N E N K H G T R C A G E V A A A A N N S H C T V G I A F N A K

ATCGGAGGAGTGCGAATGCTGGACGGAGATGTCAACGACATGGTTGAAGCAAAATCAGTTAGCTTCAACCCCGCAGCAGTGCAACATTTACAGCGCCAGCTGGGCGCGGATGATGATGC 840
I G G V R M L D G D V T D M V E A K S V S F N P Q H V H I Y S A S W G P D D D G

AAGACTGTGACGGACGACCGCCCTCACCGGCAAGCCTTTGAAACGGCTTAGAATGGGCGGAGAGCGCTCGGCTCTGTGTTTGTGTCATCTGGAATGTTGGAAGGAGCAAA 960
K T V D G P A P L T R Q A F E N G V R M G R G L G S V F V W A S G N G G R S K

GACCACTGCTCCTGTGATGCTACCAACAGCATCTACACCATCTCCATCAGCAGCAGCTGCAGAAAGCGGAAGAAACCTTGTAAGTGAAGAGTGTTCATCCACGCTGCCACAAC 1080
D H C S C D G Y T N S I Y T I S I S S T A E S G K K P W Y L E C S S T L A T T

TACAGACGGGGAGTCTACGATAAGAAATCATCACTACAGATCTGAGGCAGGTTGCACGACAAACCACTGGAGCTGAGCCTCAGCCCCCATGGCTGCAGGCATCATTCGGCTG 1140
Y S S G E S Y D K K I I T T D L R Q R C T D N H T G T S A S A P M A A G I I A L

GCCCTGGAAGCCAATCCGTTCTGACCTGGAGAGAGGTACAGCATGTTATTGTACGACTTCCCGTGGGACATTTGAACGCTAATGACTGGAAACCAATGCTGCTGCTTTAAGGTG 1320
A L E A N P F L T W R D V Q H V I V R T S R A G H L N A N D W K T N A A G F K V

AGCCATCTTTATGGATTTGGACTGATGGACGCAAGCCATGGTGATGGAGGACAGAGTGGACCAACGTTCCCGCGCAGCAGCTGTGTGTGGAGAGCACAGCCGACAAATCAAGACA 1440
S H L Y G F G L M D A E A M V M E A E K W T T V P R Q H V C V E S T D R Q I K T

2/26

FILE - 2A

3/26

ATCCGCCCTAACAGTGCAGTGGCTCCATCTACAAAGCTTCAGGCTGCTCGGATAACCCCAACGCCCATGTCAACTACCTGGAGCAGCTCGTTGTGGCATCACCATCACCACCCAGG 1560
 I R P N S A V R S I Y K A S G C S D N P N R H V N Y L E H V V V R I T I T H P R
 AGAGAGACCTGGCCATCTACCTGACCTCGCCCTCTGGAACCTAGGTCTCAGCTTTTGGCCACAGGCTATTTGATCACTCCATGGAAGGATTCAAAAACCTGGAGTTTCATGACCATTCTAT 1680
 R G D L A I Y L T S P S G T R S Q L L A N R L F D H S M E G F K N W E F M T I H
 TGCTGGGAGAAAGAGCTGCTGGTGACTGGGTCTTGAAGTTTATGATATCTCCCTCTCAGCTAAGGAACTTTAAGACTCCAGGTAAATTGAAGAATGGTCTTTGGTCTCTACGGCACC 1800
 C W G E R A A A G D W V L E V Y D T P S Q L R N F K T P G K L K E W S L V L Y G T
 TCCGTGCAGCCATATTACCAACCAATGAATTCGGAAGTGGAAAGGTTCCGCTATAGCCGAGTTGAAGACCCACAGACGACTATGSCACAGAGGATTATGCAGGTCCCTGCCACCCT 1920
 S V Q P Y S P T N E F P K V E R F R Y S R V E D P T D D Y G T E D Y A G P C D P
 GAGTGCAGTGAGGTGGCTGTGACGGGCCAGGACAGACCACTGCAATGACTGTTTGCACACTACTACTACAAGCTGAAACAAATCCAGGATCTGTGTCTCCAGCTGCCCCCTGCCAC 2040
 E C S E V G C D G P G P D H C N D C L H Y Y K L K N N T R I C V S S C P P G H
 TACCACGCCGACAAGAAGCGCTGCAGGAAGTGTGCCCCCAACTGTGAGTCTGCTTTGGAGCCATGGTGACCAATGCATGTCTGCAAAATATGGATACCTTTCTGAATGAAGAAACCAAC 2160
 Y H A D K K R C R K C A P N C E S C F G S H G D Q C M S C K Y G Y F L N E E T N
 AGCTGTGTTACTCACTGCCCTGATGGGTCAATCAGGATACCAAGAAAATCTTTGCCGGAATGCAGTGAAAACCTGCAAGACATGTACTGAAATCCATACTGTACAGAATGTAGGAT 2280
 S C V T H C P D G S Y Q D T K K N L C R K C S E N C K T C T E F H N C T E C R D
 GGTAAAGCCTGCAGGATCCGGTGTCTGCTCTGGAAGATGGACGGTATTTCAAGCGCCAGGACTGCCAGCCCTGCCACCGCTTCTGCCCACTTGTGCTGGGCGAGGATCAT 2400
 G L S L Q G S R C S V S C E D G R Y F N G Q D C Q P C H R F C A T C A G A G A D
 GGTGCATTAACTGCACAGAGGCTACTTTCATGGAGATGGGAGATGCGTGCAGAGCTGTAGTATCAGCTATTACTTTGACCACTCTTCAGAGAAATGGATACAAATCCTGCAAAAATGT 2520
 G C I N C T E G Y F M E D G R C V Q S C S I S Y Y F D H S S E N G Y K S C K K C
 GATATCAGTTGTTTGACGTGCAATGGCCAGGATTCAAGAACTGTACAAGCTGCCCTAGTGGGTATCTTTAGACTTAGGAATGTGTCAAAATGGAGGCCATTTCAGAGGATGCAACGGAA 2640
 D I S C L T C N G P G F K N C T S C P S G Y L L D L G M C Q M G A I C K D A T E
 GAGTCTGGGGAGGAGGCTTCTGTATGCTTGTGAAAAGAACAACTCTGTGCCAACGGAGGTTCTTCACAACTTTGCTGCAAAACATGTACATTTCAAGGCTGAGCAGCCATCTTA 2760
 E S W A E G G F C M L V K K N N L C Q R K V L Q Q L C C K T C T F Q G STOP
 GATTCTTTGTTCTGTAGACTTATAGATTATCCATATTATTAAAGAAAAA



4/26







FIG. 3

5/26



6/26

FIG. 5

7/26




FIG. 6

8/26





Fig 7

9/26

antisense
sense
mismatch
SON
SK-N-MCIXC
PC5

FIG. 1

10/26



FIS-9

	PC1	PC2	PC5	PC7	Furin
Number of Slide	n=31	n=29	n=25	n=27	n=30
Mean Positivity (Scale 0-3)	0.29	2.03	2.08	1.19	1.08
SD	0.46	0.61	0.62	1.04	1.07
% (+) Slide	29.0	96.6	96.0	63.0	60

FIS-10

11/26

Mirandal AGCGTCGGGACCATGGATTGGGATTGGGGAACCGCTGCAGCCGCCCGG 50
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
IRCM 1ATGGACTGGGACTGGGGGAACCGCTGCAGCCGCCCGG 38

51 ACGGCGGGACCTGCTGTGCGTGCTGGCACTGCTCGCCGGCTGTCTGCTCC 100
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
39 ACGGCGGGATCTGCTGTGCGTGCTGGCGCTGCTCGGGGCTGCCTGCTCC 88

101 CGGTATGCCCGGACGCGCGTCTACACCAACCACCTGGGCAGTGAAGATCGCC 150
| | | | | ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
89 CCGTGTGTCGGACGCGCGTCTACACCAACCACCTGGGCAGTCAAAATCGCC 138

151 GGCGGCTTCGCGGAGGCAGATCGCATAGCCAGCAAGTACGGATTCAATCAA 200
| | | | | ||||| | | | | | ||||| ||||| ||||| ||||| |||||
139 GGGGCTTCCCGAGGCCAACCGTATCGCCAGCAAGTACGGATTCAATCAA 188

201 CGTAGGACAGATCGGTGCACCTGAAGACTACTATCACTTCTACCATAGTA 250
| | | | | ||||| | | | | | ||||| ||||| ||||| ||||| |

~~IIA~~ - IIA

189 CATAGGACAGATAGGGCCCTGAAGGACTACTACCACTTCTACCATAGCA 238
251 GGACCATTAAGGTCTGTTCTCTCGAGCAGAGGAACCCACAGTTTCATT 300
239 GGACGATTAAAGGTCAGTTATCTCGAGCAGAGGGACCCACAGTTTCATT 288
301 TCAATGGAAACCAAGGTGGAGTGGATCCAAACAGCAAGTGGTGAAAAAAG 350
289 TCAATGGAAACCAAGGTGGAATGGATCCAAACAGCAAGTGGTAAAAAAGCG 338
351 AACCAAGAGGGATTATGACCTCAGCCCATGCCCAGTCAACCTACTTCAATG 400
339 GACAAAGAGGGATTATGACTTCAGTCTGTCGCCCAGTCTACCTATTTC AATG 388
401 ATCCCAAGTGGCCCAAGTATGTGGTACATGCACCTGTAGCGACAATACACAT 450
389 ATCCCAAGTGGCCCAAGCATGTGGTATATGCACCTGCAGTGACAATACACAT 438
451 CCCTGCCAGTCTGACATGAATATCGAAGGAGGCCTGGAAGAGAGGCTACAC 500

File: 11B

13/26

439 CCTGCCAGTCTGACATGAATATCGAAGGAGCCTGGAAGAGAGGCTACAC 488
501 GGGAAAGAACATTTGTGGTCACTATCCTGGATGACGGAATTGAGAGAACCC 550
|||||
489 GGGAAAGAACATTTGTGGTCACTATCCTGGATGACGGAATTGAGAGAACCC 538
551 ATCCAGATCTGATGC AAAA C TACGATGCTCTGGCAAAGTTGCCGACGTGAAT 600
|||||
539 ATCCAGATCTGATGC AAAA C TACGATGCTCTGGCAAAGTTGCCGACGTGAAT 588
601 GGGAAATGACTTGGACCCCAATGCCCTCGTTATGATGCAAGCAACGAGAACAA 650
|||||
589 GGGAAATGACTTGGACCCCAATGCCCTCGTTATGATGCAAGCAACGAGAACAA 638
651 GCATGGGACTCGCTGTGCTGGAGAAAGTGGCAGCCGCTGC AAACAATTTCGC 700
|||||
639 GCATGGGACTCGCTGTGCTGGAGAAAGTGGCAGCCGCTGC AAACAATTTCGC 688
701 ACTGCACAGTCGGAATTGCTTTCAACGCCCAAGATCGGAGGAGTGCGAATG 750
|||||

Ex - 11C

14/26

689 ACTGCACAGTCGGAATTGCTTTCAACGCCAAGATCGGAGGAGTGCGAATG 738
751 CTGGACGGAGATGTCACGGACATGGTTGAAGCAAAATCAGTTAGCTTCAA 800
739 CTGGACGGAGATGTCACGGACATGGTTGAAGCAAAATCAGTTAGCTTCAA 788
801 CCCCAGCACGTGCACATTACAGCGCCAGCTGGGGCCCGGATGATGATG 850
789 CCCCAGCACGTGCACATTACAGCGCCAGCTGGGGCCCGGATGATGATG 838
851 GCAAGACTGTGGACGGACCAAGCCCCCTCACCCGGCAAGCCTTTGAAAAC 900
839 GCAAGACTGTGGACGGACCAAGCCCCCTCACCCGGCAAGCCTTTGAAAAC 888
901 GGCGTTAGAAATGGGGCGGAGAGGCCCTCGGCTCTGTGTTTGGGCATC 950
889 GGCGTTAGAAATGGGGCGGAGAGGCCCTCGGCTCTGTGTTTGGGCATC 938
951 TGGAAATGGTGAAGGAGCAAGACCACTGCTCCTGTGATGGCTACACCA 1000

Ex - 11D

15/26

939 TGGAAATGGTGGAAGGAGCAAAAGACCACTGCTCCTGTGATGGCTACACCA 988
1001 ACAGCATCTACACCATCTCCATCAGCAGCACTGCAGAAAGCGGAAAGAAA 1050
|||||
989 ACAGCATCTACACCATCTCCATCAGCAGCACTGCAGAAAGCGGAAAGAAA 1038
1051 CCTTGGTACCTGGAAGAGTGTTCCATCCACGCTGGCCACAACCTACAGCAG 1100
|||||
1039 CCTTGGTACCTGGAAGAGTGTTCCATCCACGCTGGCCACAACCTACAGCAG 1088
1101 CGGGAGTCCCTACGATAAGAAAATCATCACTACAGATCTGAGGCAGCGTT 1150
|||||
1089 CGGGAGTCCCTACGATAAGAAAATCATCACTACAGATCTGAGGCAGCGTT 1138
1151 GCACGGACAACCACTGGGACGTCAGCCTCAGCCCCCATGGCTGCAGGC 1200
|||||
1139 GCACGGACAACCACTGGGACGTCAGCCTCAGCCCCCATGGCTGCAGGC 1188
1201 ATCATTCGCGCTGGCCCTGGAAAGCCAAATCCGTTTCTGACCTGGAGAGACGT 1250
|||||

FILE

16/26

1189 ATCATTCGCGCTGGCCCTGGAAGCCAATCCGTTTCTGACCTGGAGAGACGT 1238
1251 ACAGCATGTTATTGTCAGGACTTCCCGTGCGGGACATTTGAACGCTAATG 1300
|||||
1239 ACAGCATGTTATTGTCAGGACTTCCCGTGCGGGACATTTGAACGCTAATG 1288
1301 ACTGGAAACCAATGCTGCTGGTTTAAAGGTGAGCCATCTTTATGGATT 1350
|||||
1289 ACTGGAAACCAATGCTGCTGGTTTAAAGGTGAGCCATCTTTATGGATT 1338
1351 GGACTGATGGACGCAGAAAGCCATGGTGTGATGGAGGCAGAGAAGTGGACCAC 1400
|||||
1339 GGACTGATGGACGCAGAAAGCCATGGTGTGATGGAGGCAGAGAAGTGGACCAC 1388
1401 CGTTCCTCCCGCAGCACGTGTGTGTGGAGAGCACAGACCCGACAAATCAAGA 1450
|||||
1389 CGTTCCTCCCGCAGCACGTGTGTGTGGAGAGCACAGACCCGACAAATCAAGA 1438
1451 CAATCCGCCCTAACAGTGCAGTGCCTCCATCTACAAAGCTTCAGGCTGC 1500
|||||

FEF - 11F

17/26

1439 CAATCCGCCCTAACAGTGCAGTGCCTCCATCTACAAAGCTTCAGGCTGC 1488
1501 TCGGATAACCCCAACCGCCATGTCAACTACCTGGAGCACGTCGTTGTGCG 1550
|||||
1489 TCGGATAACCCCAACCGCCATGTCAACTACCTGGAGCACGTCGTTGTGCG 1538
1551 CATCACCATCACCCACCCAGGAGAGAGACCTGGCCATCTACCTGACCT 1600
|||||
1539 CATCACCATCACCCACCCAGGAGAGAGACCTGGCCATCTACCTGACCT 1588
1601 CGCCCTCTGGAACTAGGTCTCAGCTTTGGCCAAACAGGCTATTGATCAC 1650
|||||
1589 CGCCCTCTGGAACTAGGTCTCAGCTTTGGCCAAACAGGCTATTGATCAC 1638
1651 TCCATGGAAGGATTCAAAAACCTGGGAGTTCATGACCATTTCATTGCTGGGG 1700
|||||
1639 TCCATGGAAGGATTCAAAAACCTGGGAGTTCATGACCATTTCATTGCTGGGG 1688
1701 AGAAAGAGCTGCTGGTGACTGGGTCCCTTGAAGTTTATGATACTCCCTCTC 1750
|||||

- 116

18/26

1689 AGAAAGAGCTGCTGGTGACTGGGTCCTTGAAGTTATGATACTCCCTCTC 1738
1751 AGCTAAGGAACCTTAAAGACTCCAGGTAAATTGAAAGAAATGGTCTTTGGTC 1800
|||||
1739 AGCTAAGGAACCTTAAAGACTCCAGGTAAATTGAAAGAAATGGTCTTTGGTC 1788
1801 CTCACGGCACCTCCGTCCGGGCCATATTCACCAACCAATGAATTTCCGAA 1850
|||||
1789 CTCACGGCACCTCCGTCCAGGCCATATTCACCAACCAATGAATTTCCGAA 1838
1851 AGTGGAACGGTTCCGCTATAGCCGAGTTGAAGACCCACAGACGACTATG 1900
|||||
1839 AGTGGAACGGTTCCGCTATAGCCGAGTTGAAGACCCACAGACGACTATG 1888
1901 GCACAGAGGATTATGCAGGTCCCTGCGACCCCTGAGTGCAGTGAGGTGGC 1950
|||||
1889 GCACAGAGGATTATGCAGGTCCCTGCGACCCCTGAGTGCAGTGAGGTGGC 1938
1951 TGTGACGGGCCAGGACCAGACCACTGCAATGACTGTTGCACTACTACTA 2000
|||||

FILED - 11H

19/26

1939 TGTGACGGCCAGGACCAGACCAGCAATGACTGTTGCACTACTACTA 1988
2001 CAAGCTGAAACAAATACCAAGGATCTGTGTCTCCAGCTGCCCCCCCTGGCC 2050
|||||
1989 CAAGCTGAAACAAATACCAAGGATCTGTGTCTCCAGCTGCCCCCCCTGGCC 2038
2051 ACTACCACGCCGACAAGACGCTGCAGGAAGTGTGCCCCCAACTGTGAG 2100
|||||
2039 ACTACCACGCCGACAAGACGCTGCAGGAAGTGTGCCCCCAACTGTGAG 2088
2101 TCCTGCTTTGGGAGCCATGGTGACCAATGCATGTCTGCAAAATATGGATA 2150
|||||
2089 TCCTGCTTTGGGAGCCATGGTGACCAATGCATGTCTGCAAAATATGGATA 2138
2151 CTTTCTGAATGAAGAAACCAACAGCTGTGTACTCACTGCCCTGATGGGT 2200
|||||
2139 CTTTCTGAATGAAGAAACCAACAGCTGTGTACTCACTGCCCTGATGGGT 2188
2201 CATATCAGGATACCAAGAAATCTTTGCCGAAATGCAGTGAAACTGC 2250
|||||

FIG - III

20/26

2189 CATATCAGGATACCAAGAAAAATCTTTGCCGGAATGCAGTGAAACTGC 2238
2251 AAGACATGTA[.]CTGAATTC[.]CATAA[.]CTGTACAGAATGTAGGATGGGTTAAG 2300
|||||
2239 AAGACATGTA[.]CTGAATTC[.]CATAA[.]CTGTACAGAATGTAGGATGGGTTAAG 2288
2301 CCTGCAGGATCCC[.]GGTGCTCTGTCTCCTCTGTGAAGATGGACGGTATTCA 2350
|||||
2289 CCTGCAGGATCCC[.]GGTGCTCTGTCTCCTCTGTGAAGATGGACGGTATTCA 2338
2351 ACGGCCAGGACTGCCAGCCCTGCCACCGCTTCTGCGCCACTTGTGCTGGG 2400
|||||
2339 ACGGCCAGGACTGCCAGCCCTGCCACCGCTTCTGCGCCACTTGTGCTGGG 2388
2401 GCAGGAGCTGATGGGTGCATTA[.]ACTGCACAGAGGGCTACTTCATGGAGGA 2450
|||||
2389 GCAGGAGCTGATGGGTGCATTA[.]ACTGCACAGAGGGCTACTTCATGGAGGA 2438
2451 TGGGAGATGCCGTGCAGAGCTGTAGTATCAGCTATTACTTTGACCACTCTT 2500
|||||

FIG - 11J

21/26

2439 TGGGAGATGCCGTGCAGAGCTGTAGTATCAGCTATTACTTTGACCACCTCTT 2488
2501 CAGAGAAATGGATACAAATCCCTGC AAAAATGTGATATCAGTTGTTGACG 2550
|||||
2489 CAGAGAAATGGATACAAATCCCTGC AAAAATGTGATATCAGTTGTTGACG 2538
2551 TGCAATGGCC CAGGATTCAAGAACTGTACAAGCTGCCCTAGTGGGTATCT 2600
|||||
2539 TGCAATGGCC CAGGATTCAAGAACTGTACAAGCTGCCCTAGTGGGTATCT 2588
2601 CTTAGACTTAGGAATGTGTCAAAATGGGAGCCATTGCAAGGATGCAACGG 2650
|||||
2589 CTTAGACTTAGGAATGTGTCAAAATGGGAGCCATTGCAAGGATGCAACGG 2638
2651 AAGAGTCCTGGGCGGAAGGAGGCTTCTGTATGCTTGTGAAAAGAACAAAT 2700
|||||
2639 AAGAGTCCTGGGCGGAAGGAGGCTTCTGTATGCTTGTGAAAAGAACAAAT 2688
2701 CTGTGCCAACGGAAGGTTCTTCAACAACTTTGCTGCAAAACATGTACATT 2750
|||||

FEI - 11K

22/26

2689 CTGTGCCAACGGAAGGTTCTTCAACAACCTTGTCTGCAAAACATGTACATT 2738

2751 CCAAGGCTGAGCAGCC..... 2766

| | | | | | | | | |

2739 TCAAGGCTGAGCAGCCATCTTAGATTCTTTGTTCCCTGTAGACTTATAGA 2788

. . .

Figure - 111

23/26

Miranda1	MDWDWGNRCSRPGRRDLLCVLALLAGCLLPVCRTRVYTNHWAVKIAGGFA	50
IRCM 1	MDWDWGNRCSRPGRRDLLCVLALLGGCLLPVCRTRVYTNHWAVKIAGGFP	50
51	EADRIASKYGFINVGGQIGALKDYHFFYHSRTIKRSVLSSRGTHSFISMEP	100
	. : : : : :	
51	EANRIASKYGFINIGQIGALKDYHFFYHSRTIKRSVISSRGTHSFISMEP	100
101	KVEWIIQQQVVKRTRKRDYDLSHAQSTYFNDEPKWPSMWYMHCSDNTHPCQS	150
101	KVEWIIQQQVVKRTRKRDYDFSRAQSTYFNDEPKWPSMWYMHCSDNTHPCQS	150
151	DMNIEGAWKRGYTGKNI VVTILDDGIERTHPDLMQNYDALASCDVNGNDL	200
151	DMNIEGAWKRGYTGKNI VVTILDDGIERTHPDLMQNYDALASCDVNGNDL	200
201	DPMPRYDASNENKHGTRCAGEVAAAANNHCTVGIAFNAKIGGVRLDGD	250
201	DPMPRYDASNENKHGTRCAGEVAAAANNHCTVGIAFNAKIGGVRLDGD	250

712A

24/26

251 VTDMVEAKSVSNPQHVIYSASWGPDGKTVDGPAPLTRQAFENGVRM 300
 |||||
 251 VTDMVEAKSVSNPQHVIYSASWGPDGKTVDGPAPLTRQAFENGVRM 300
 |||||
 301 GRRGLGSVFVWASNGGRSKDHCSCDGYTNSIYTISSSTAESGKKPWYL 350
 |||||
 301 GRRGLGSVFVWASNGGRSKDHCSCDGYTNSIYTISSSTAESGKKPWYL 350
 |||||
 351 EECSSLATYSSGESYDKKIIITDLRQRCTDNHTGTSASAPMAAGIIAL 400
 |||||
 351 EECSSLATYSSGESYDKKIIITDLRQRCTDNHTGTSASAPMAAGIIAL 400
 |||||
 401 ALEANPFLTWRDVQHVIVRTSRAGHLNANDWKTNAAGFKVSHLYGFGGLMD 450
 |||||
 401 ALEANPFLTWRDVQHVIVRTSRAGHLNANDWKTNAAGFKVSHLYGFGGLMD 450
 |||||
 451 AEAMVMEAEKWTTPRQHVCVESTDRQIKTIRPNSAVRSIYKASGCSDNP 500
 |||||
 451 AEAMVMEAEKWTTPRQHVCVESTDRQIKTIRPNSAVRSIYKASGCSDNP 500

128

26/26

[illegible]

027-5111



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/64, 15/11, 15/57		A3	(11) International Publication Number: WO 98/04686 (43) International Publication Date: 5 February 1998 (05.02.98)
(21) International Application Number: PCT/CA97/00535 (22) International Filing Date: 25 July 1997 (25.07.97) (30) Priority Data: 60/021,008 26 July 1996 (26.07.96) US 2,203,745 25 April 1997 (25.04.97) CA (71)(72) Applicants and Inventors: DAY, Robert [CA/CA]; 833 rue Pierre, Sainte-Dorothée, Québec H2W 3S6 (CA). SEIDAH, Nabil, G. [CA/CA]; Apartment 1412, 200 de Gaspé, Îles-des-Sœurs, Québec H3E 1S6 (CA). MARTEL, Rémi [CA/CA]; 4865 Lafontaine, Montréal, Québec H1V 1R7 (CA). CHRETIEN, Michel [CA/CA]; Apartment 1404, 1 Côte Sainte-Catherine, Montréal, Québec H2V 1Z8 (CA). REUDELHUBER, Tim [CA/CA]; 671 Warwick Drive, Baie d'Urfé, Québec H9X 2P4 (CA). LECLERC, Guy [CA/CA]; 327 Lorraine, Rosemère, Québec J7A 4K1 (CA). (74) Agents: DUBUC, Jean, H. et al.; Goudreau Gage Dubuc & Martineau Walker, 800 Place Victoria Tower, Suite 3400, P.O. Box 242, Montreal, Quebec H4Z 1E9 (CA).			(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 23 April 1998 (23.04.98)
(54) Title: PRO-PROTEIN CONVERTING ENZYME			
(57) Abstract			
<p>A cDNA clone encoding the human prohormone convertase PC5 was isolated from human adrenal gland messenger RNA. The deduced protein sequence would encode a 915 amino acid prepro PC5 which shares a very high degree of homology with the previously cloned rat and mouse homologues. PC5 mRNA is detected in multiple human tissues, including the brain, adrenal and thyroid glands, heart, placenta, lung and testes. PC5 mRNA was undetectable in the liver and is present at lower levels in skeletal muscle, kidney, pancreas, small intestine and stomach. Co-transfection of human PC5 and human prorenin expression vectors in cultured GH4Cl cells leads to secretion of active renin. The activation of human prorenin by PC5 is dependent on a pair of basic amino acids at positions 42 and 43 of the prorenin prosegment and occurs only in cells containing dense core secretory granules. Human PC5 was co-localized with renin by immunohistochemistry in the zona glomerulosa of the adrenal gland suggesting that it could participate in the activation of a local renin-angiotensin system in the human adrenal cortex. PC5 is overexpressed in atherosclerotic coronary blood vessels. Silencing PC5 expression with a specific antisense oligonucleotide efficiently inhibited the proliferation of smooth muscle cells in culture. Furthermore, the antisense inhibited carotid stenosis in a carotid injury model. These results indicate that silencing PC5 applies to the prevention of restenosis. PCs could be targets of choice for treating any proliferative diseases involving their action on a given growth factor. Finally, the antisense oligonucleotide PC5 is to be used for silencing the activity of this enzyme towards HIV gp160, since both coexist in CD4⁺ T lymphocytes and the viral glycoprotein is cleavable by PC5.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 97/00535

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/64 C12N15/11 C12N15/57

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	MERCURE C. ET AL.: "Prohormone convertase PC5 is a candidate processing enzyme for prorenin in the human adrenal cortex" HYPERTENSION, vol. 28, no. 4, October 1996, DALLAS, US, pages 840-846, XP002046402 see the whole document --- -/-	1-5

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

17 November 1997

Date of mailing of the international search report

10.03.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Panzica, G

INTERNATIONAL SEARCH REPORT

International Application No
PCT/CA 97/00535

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MIRANDA L. ET AL.: "Isolation of the human PC6 gene encoding the putative host protease for HIV-1 gp160 processing in CD4+ T lymphocytes" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, July 1996, WASHINGTON US, pages 7695-7700, XP002046403 cited in the application	1,3-6,8, 12
Y	see the whole document	11, 13-15,18
Y	--- WO 94 15945 A (TEXAS BIOTECHNOLOGY CORPORATION) 21 July 1994 see abstract see page 2, line 10 - line 22 ---	11, 13-15,18
A	--- DECROLY E ET AL: "THE CONVERTASES FURIN AND PC1 CAN BOTH CLEAVE THE HUMAN IMMUNODEFICIENCY VIRUS (HIV)-1 ENVELOPE GLYCOPROTEIN GP160 INTO GP120 (HIV-I SU) AND GP41 (HIV-I TM)" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 269, no. 16, 22 April 1994, pages 12240-12247, XP002023318 ---	8,12-18
A	--- HALÈNE C ET AL: "SPECIFIC REGULATION OF GENE EXPRESSION BY ANTISENSE, SENSE AND ANTIGENE NUCLEIC ACIDS" BIOCHIMICA ET BIOPHYSICA ACTA, vol. 1049, 1 January 1990, pages 99-125, XP000570355 -----	

INTERNATIONAL SEARCH REPORT

Inter. application No.
PCT/CA 97/00535

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1 8 11 15
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
The definition of an enzyme by the term PC5 is not sufficient to define the subject matter as it is a non structural definition. The search has been performed according to the definition of the subject matter as given by the Sequences Nos. 1 and 2 of the seq. listing.
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/CA 97/00535

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-6, 8, 11-15, part. 16-18

A purified human pro-protein converting enzyme isolated from adrenals, its DNA and aminoacid sequences and its uses.

2. Claims: 7, 9, 10, part. 16-18

An antisense oligonucleotide and its uses

INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/CA 97/00535

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9415945 A	21-07-1994	AU 6080194 A	15-08-1994
		CA 2152903 A	21-07-1994
		EP 0677055 A	18-10-1995
